MIXED ESTERS OF LACTIC AND CARBONIC ACIDS. n-ALKYL CARBONATES OF n-ALKYL LACTATES

Previous papers in this series described a group of miscellaneous carbonates of lactates (1) and three homologous families (2), and also indicated the usefulness of some of these esters as plasticizers for vinyl chloride resins (3). This paper describes two additional series of esters having the formula ROCOOCH $(CH_3)COOR'$. In one homologous series, R is ethyl and R' is n-alkyl. In the other, R and R' are identical n-alkyl groups. Each series consists of six members, one compound being common to both (Table I).

The esters were prepared, and physical properties were determined as described in a previous paper (2). The preparation and physical properties of the lactates used have been described recently (4).

Although the lowest members of each series have been prepared (2), they are not included in this paper because, as is usual in homologous series, their physical properties deviate from the orderly sequence shown by the higher members of the series.

Boiling points and vapor pressures. Figures 1 and 2 show the boiling points of the esters as a function of the pressure. In these modified Cox charts, pressure is plotted logarithmically, and the temperature scale is laid off as a linear function of 1/(t+193), where t is in °C. The lines showed considerable curvature (convex upward) when plotted on the usual types of Cox paper having temperature scales determined by 1/(t+273) or 1/(t+230), although the curvature was less with the latter type paper. The scales determined by 1/(t+193) were straight lines, and charts having this scale were conveniently prepared from commercial Cox chart paper having a scale linear with 1/(t+273) by adding 80° to each temperature designated on the chart. The mathematical proof of this transformation has been published elsewhere (5).

In each homologous series, the logarithm of the vapor pressure at any fixed temperature is a linear function of the number of carbon atoms in the ester. Tables II and III show equations for these lines at various temperatures. For each series, these lines constitute a family having a common point of intersection located as follows:

For the ethyl carbonate series: x = -18, $\log P = 7.5$.

For the alkyl carbonate series: x = -14.5, $\log P = 6.5$.

Also, within each family, the slopes (a) of these lines vary with the absolute temperature:

For the ethyl carbonate series: a = -113/T + 0.055.

For the alkyl carbonate series: a = -130/T + 0.103.

¹ One of the Laboratories of the Bureau of Agricultural and Industrial Chemistry, Agricultural Research Administration, United States Department of Agriculture.

TABLE I

n-Alkyl Carbonates of n-Alkyl Lactates: ROCOOCHICH M

	Н,%	-	round								9.5			
CHARLES OF W-ALKYL LACTATES: ROCOOCH(CH1)COOR,		1	Carca	7.4	8.3	9.0	9.6	10.4	10.9	8.3	0.6	9.6	10.0	11.9
	C, %	Found									59.1			
	0	Calc'd		50.5	55.0	58.5	61.3	65.4	68.4	55.0	58.5	61.3	63.5	69.5
	VISCOSITY, CPS.	400		2.45	2.92	3.79	4.61	7.51	11.57	8.8	8. 24.	4.57	5.65	11.68
	VISCOSI	20°		4.11	5.16	7.46	8.96	15.64	25.48	2.88	6.51	8.74	11.59	25.66
	MOL, REFR. 20°	Found		43.98	53.28	62.75	71.70	90.41	108.97	93.31	02.70	70.17	91.04	118.27
		Calc'd		44.10	53.33	27.20	2 5	100.27	100.70	80.00	70.20	200	117.04	11/.98
	d.40		0000	1.0523	1.0138	0.3020	0.000	0.036	1 0138	0.000	0.900	0.000	0.010	O. Stor
	d.20		1 07.10	1.0/42	1.0021	0 0846	0 0560	0 0300	1 0338	1 0040	0.9848	0.9654	0 0315	- 0.00.0
	240 D		1 4090	1.1002	1 4167	1,4212	1.4295	1.4352	1.4110	1.4162	1.4214	1.4250	1.4358	-
	, 20 0		1 4119	1.4181	1.4242	1.4281	1.4366	1.4422	1.4189	1.4240	1.4290	1.4327	1.4440	
	VIELD, %		72	' Z	8	49	48	8	22	23	29	49	23	
	ĸ		Ethyl	Butyl	Hexyl	Octylb	Dodecyl	Hexadecyl	Propyl	Butyla	Amyle	Hexyle	Decyl•	
	æ		Ethyl	Ethyl	Ethyl	Ethyl	Ethyl	Ethyl	Propyl	Butyl	Amyl	Hexyl	Decyl	

^a These compounds have been described previously (2), but are included here for comparison with their homologs. ^b Some of the data on this compound have been published previously (1). ^c Certain data on these compounds have been reported previously (3).

By use of these equations for the slopes, and the common points given above, equations for vapor pressures at any desired temperature may be readily calculated.

At any chosen fixed pressure, the squares of the boiling points (°K) vary directly with the number of carbon atoms in the esters of either series. Equations in Tables II and III show these relationships. Within each family, these lines determined at different pressures pass through a common point.

TABLE II

BOILING POINTS (T = $^{\circ}$ K.) AND VAPOR PRESSURES (P = PRESSURE, MM.) OF ETHYL

CARBONATES OF n-ALKYL LACTATES AS RELATED TO NUMBER OF

CARBON ATOMS (x)

•C			DEVIATIONS ⁴			
CONST. TEMP., °C., or PRESSURE, MM.	а	b	Max.	Average		
and the second s	Boiling	Points: 10 ⁻⁴ T ² =	ax + b			
760 mm.	1.56	11.30	3	1.2		
100	1.18	8.50	1	0.7		
10	0.918	6.50	3	1.8		
1	0.746	5.18	4	2.0		
	Vapor P	ressures: Log P =	ax + b			
300°	-0.143	4.92	19	14		
250	-0.161	4.57	9	7		
200	-0.184	4.17	6	2		
150	-0.212	3.67	2	1		
100	-0.248	3.00	6	4		

^a Temperature deviations are in °C. Pressure deviations are in percentages. A deviation of 1° is equivalent to a pressure deviation of about 5%.

TABLE III

BOILING POINTS (T = $^{\circ}$ K.) AND VAPOR PRESSURES (P = PRESSURE, MM.) OF COMPOUNDS ROCOOCH(CH₂)COOR (R = n-Alkyl) as Related to Number OF Carbon Atoms (x)

• •	а		deviations ^a			
CONSTANT TEMP., °C. or pressure, mm.		b	Max.	Average		
	Boiling	Points: 10-4T2 =	ax + b			
760 mm.	1.51	11.95	2	0.6		
100	1.15	8.80	5	1.7		
10	0.896	6.60	2	1.0		
1	0.734	5.12	2	1.3		
	Vapor P	ressures: Log P	= ax + b			
300°	-0.127	4.66	11	7		
250	-0.145	4.35	7	4		
200	-0.171	4.02	3	2		
150	-0.202	3.57	4	2		
100	-0.245	2.95	4	3		

^a Temperature deviations are in °C. Pressure deviations are in percentages. A deviation of 1° is equivalent to a pressure deviation of about 5%.

For ethyl carbonates: x = -7.5, $10^{-4}T^2 = -0.4$. For alkyl carbonates: x = -9.0, $10^{-4}T^2 = -1.5$. The slopes (a) of these lines vary inversely with the logarithm of the pressure

For the ethyl carbonates: Log P = -4.21/a + 5.59.

For the alkyl carbonates: Log P = -4.21/a + 5.69.

By use of these equations for the slopes and the common points given above, equations may be calculated for boiling points at any desired pressure.

Densities, refractive indices, and viscosities. These physical properties were measured at 20° and at 40° (Table I). As reported in the previous paper (2), linear relationships were found between certain functions of these physical constants and the number of carbon atoms in the esters. Table IV shows equations for these relationships. These equations are highly useful, not only for calculating the properties of homologs not prepared but for checking the purity

TABLE IV Equations Relating Refractive Indices, Densities, and Viscosities to the Number OF CARBON ATOMS IN CARBONATES OF LACTATES ROCOOCH(CH3)COOR'

OF CAR	CBON ATOMS	IN CARBONATES OF LACTATES ROCO	DEVIATIONS		
R	R'	EQUATION	Max.	Average	
Ethyl Ethyl Ethyl Ethyl Ethyl Ethyl R = R' =	n-Alkyl	$1/(x+10) = -0.773 n_D^{20} + 1.1462$ $1/(x+10) = -0.741 n_D^{40} + 1.095$ $1/x = 0.583 d_A^{40} - 0.5019$ $1/x = 0.618 d_A^{4} - 0.5253$ $Log \eta_{20} = 0.0566 x + 0.161$ $Log \eta_{40} = 0.0495 x - 0.030$ $1/(x+8) = -0.954 n_D^{20} + 1.4012$ $1/(x+8) = -0.954 n_D^{40} + 1.4012$ $1/(x+1) = 0.498 d_A^{20} - 0.4238$ $1/(x+1) = 0.521 d_A^{40} - 0.4371$ $Log \eta_{20} = 0.0500 x + 0.240$ $Log \eta_{40} = 0.0424 x + 0.065$	0.0007 0.0005 0.0017 0.0020 6.7 ^b 3.3 ^b 0.0005 0.0005 0.0005 0.0002 ^c 7.3 ^b 3.8 ^b	$\begin{array}{c} 0.0002 \\ 0.0002 \\ 0.0008 \\ 0.0009 \\ 2.3^{b} \\ 1.6^{b} \\ 0.0002 \\ 0.0002 \\ 0.0002^{c} \\ 0.0002^{c} \\ 5.4^{b} \\ 2.6^{b} \end{array}$	

[•] Difference between calculated and observed values of the physical constants; methyl esters excluded. b Percentage deviation from the observed value. Hexyl carbonate of hexyl lactate excluded; its deviation was .0037 at 20° and .0035 at 40°.

of those studied and the accuracy of physical measurements made on them. When such correlations of physical properties with molecular structure have been made for a sufficient number of families of compounds, broader and more fundamental relationships may become demonstrable.

Acknowledgment. The authors are grateful to C. O. Willits, C. L. Ogg, and their associates for analyses, and to H. L. Fisher, U. S. Industrial Chemicals Corporation, for hexyl and decyl chloroformates.

SUMMARY

Two homologous series of n-alkyl carbonates of n-alkyl lactates, ROCOOCH (CH₃)COOR', were prepared, and several physical properties were determined. In one series R is ethyl and R' is n-alkyl; in the other R and R' are identical n-alkyl groups.

Equations were developed which relate vapor pressures, boiling points, refractive indices, densities, and viscosities to the number of carbon atoms in the members of each series.

PHILADELPHIA 18, PA.

REFERENCES

- (1) REHBERG AND DIXON, J. Org. Chem., 13, 254 (1948).
- (2) REHBERG AND DIXON, J. Org. Chem., 15, 565 (1950).
- (3) Rehberg, Dixon, and Meiss, Plasticizers from Lactic Acid. Alkyl Carbonates of Lactates, presented before the Division of Paint, Varnish, and Plastics Chemistry, American Chemical Society, Washington Meeting, August-September 1948.
- (4) REHBERG AND DIXON, J. Am. Chem. Soc., 72, 1918 (1950).
- (5) REHBERG, Ind. Eng. Chem., 42, 829 (1950).